



Spider Mites—Botanical Control Formulations

Spider mites are among the most notorious of pests. They multiply rapidly in hot, dry conditions and can easily overwhelm a field, particularly if plants are water stressed, or a greenhouse in weeks or even days. They have been found on almost 4,000 different host species (Attia et al., 2013).

Essential Oils

Spider mites are especially vulnerable to various essential oils, but so are some crops. Essential oils are distilled or cold pressed from plants. These oils do not persist in the environment like synthetic chemicals. Essential oils found to be highly effective against spider mites include garlic, white mustard, epazote, English lavender, basil, rosemary, mint, lemon balm, tansy, pennyroyal, wormwood, neem, true myrtle, orange, wild tomato, stinging nettle, and others (Mozaffaria et al., 2012).

Essential oils have toxic effects on some crops and vary in composition and potency, depending on variables such as climate, variety, and timing of harvest.

Dilution rate per gallon of water should be 10 ml to 40 ml, depending on the plant species and oils being used. If you are unsure about phytotoxicity on a specific crop, try a small test area or test plant before fully committing, especially during hot weather.



Spider mites can build up so rapidly under the right conditions that they can completely cover crops in red. Photo: Juan Raygoza

Cilantro

Using essential oil or crude extracts, and even interplanting cilantro with other crops, can prevent and kill spider mites. One simple preparation is 200 grams of crushed seeds boiled in one liter of water for 10 minutes. This extract is then strained to remove the seeds and applied to affected plants, ensuring that the undersides of the leaves are thoroughly covered (Abo-Shanab et al., 2013).

Aloe Vera

Spider mites can be smothered by the surfactant qualities of aloe vera juice. It can be mixed with neem, spinosad, or essential oils. One recipe calls for one tablespoon of neem oil, ¼ teaspoon of a silica product like Sodo Hydro, and ¼ cup of aloe juice. This solution should be sprayed thoroughly over the entire plant every three days to break up the reproductive cycle. A group of scientists also used powdered aloe leaf at a 5,000-ppm concentration in water to kill over 70% of spider mites within three days. An acetone extract caused 100% mortality in the same time period. The researchers used 100 grams of dried aloe leaf powder and 500 ml of acetone, ethyl alcohol, ethyl acetate, or water to produce the distillate. The distillate was then evaporated and dissolved into distilled water (Wei et al., 2011).

Bugleweed (*Ajuga*)

Crude extracts of bugleweed have been found to reduce spider mite populations. Crude extracts can be as simple as tossing a handful of leaves into a blender with water, straining, and then spraying judiciously onto affected plants. In Germany, researchers used a crude extract of *Ajuga remota* to effect 84% mortality of spider mites in 24 hours, with 69% mortality occurring in the first three hours. Fresh *Ajuga* juice was also shown to affect unhatched eggs and developing nymphs. The study also showed methanolic extracts of *Ajuga* being very potent against adults and highly repellent (Schauer and Schmutterer, 1981).



The photo on the left shows the characteristic webbing that is the telltale sign of spider mite infestation. The photo on the right shows field conditions that lead to spider mite epidemics: hot, dry, low humidity, and stressed plants. Photos: Rex DuFour, NCAT

Mimosa (*Albizzia julibrissin*)

Crude ethyl alcohol extracts of the common weedy mimosa tree's leaves and stems at a concentration of 10 grams per liter achieved almost 99 percent mortality of both adults and eggs of carmine spider mites. These alcohol extracts can be made at home using rum, vodka, or grain alcohol. Fill a quart-sized mason jar with freshly air-dried leaves and then pour alcohol over them, filling the jar. Close it tightly and let sit for six to eight weeks. When the infusion is complete, put about 10 ml of the liquid and one liter of distilled water in a spray bottle. Test spray one leaf to make sure the concentration is not phytotoxic. If the concentration is too high, further dilute it with distilled water. Repeat treatments as necessary (Wang et al., 2013).

Conclusion

Many plants, their crude extracts, or their essential oils can kill spider mites, but it is very important to manipulate the environment to be hostile to the spider mites. Since spider mites like hot, dry conditions, attempt to cool them down and increase humidity or even mist them, in addition to treating them with botanical pesticides.

References

Abo-Shanab, A.S.H., K.S. Moursi, and H.S. Hussein. 2013. Effect of intercropping of Coriander (*Coriandrum sativum* L.) with Tomato (*Solanum lycopersicum*) on sucking pest management infesting tomato In Nubariya, El-Bheira Governorate, Egypt. www.academia.edu/5290585/Effect_of_intercropping_of_Coriander_Coriandrum_sativum_L_with_Tomato_Solanum_lycopersicum_on_sucking_pest_management_infesting_tomato_In_Nubariya_El-Bheira_Governorate_Egypt

Attia, S., K.L. Grissa, G. Lognay, E. Bitume, T. Hance, and A. Maillieux. 2013. A review of the major biological approaches to control the worldwide pest *Tetranychus urticae* (Acari: Tetranychidae) with special reference to natural pesticides Biological approaches to control *Tetranychus urticae*. *Journal of Pest Science*. May.

Mozaffaria F, H. Abbasipoura, A.S. Garjanb, A.R. Sabooric, and M. Mahmoudvandd. 2012. Various effects of ethanolic extract of *Mentha pulegium* on the two-spotted spider mite, *Tetranychus urticae* (Tetranychidae). *Archives of Phytopathology and Plant Protection*. July. Vol. 45, No. 11. p. 1347–1355.

Schauer, M and H. Schmutterer. 1981. Wirkung von Frischpreßsäften und Rohextrakten aus der Labiate *Ajuga remota* auf die Gemeine Spinnmilbe *Tetranychus urticae* Koch. *Journal of Applied Entomology*. 91(1-5):425-433.

Wang, D.D, F.L. Jia, J. Chen , W.J. Yu , and G.H. Dai. 2013. Acaricidal Activities of *Albizzia julibrissin* Durazz Extracts against *Tetranychus cinnabarinus* (Acari: Tetranychidae). *Advanced Materials Research*. Feb. Vol. 666, pp. 77-84.

Wei, J., W. Ding, Y.-G. Zhao, P. Vanichpakorn. 2011. Acaricidal activity of *Aloe vera* L. leaf extracts against *Tetranychus cinnabarinus* Boisduval (Acarina: Tetranychidae). *Journal of Asia-Pacific Entomology*. 14 (2011) 353-356.

ATTRA Sustainable Agriculture
NCAT A program of the National Center for Appropriate Technology • 1-800-348-3140 • www.attra.ncat.org

Companion Planting & Botanical Pesticides: Concepts & Resources

By Corinne Ruppner and Matt Doshier
 Published April 2016
 Updated April 2018
 by Justin Duncan, NCAT Agriculture Specialist
 ©2018
 #1215

Contents
 Introduction 1
 Companion Planting 2
 Botanical Pesticides 3
 The Science of Companion Planting 4
 Botanical Pesticides 5
 Spider Mites 6
 Summary 7
 Appendix 8
 Glossary 9
 Bibliography 10
 Credits 11

Traditional Companion Planting
 C ompanion planting is a practice of growing different plants together to benefit them or to repel pests. It is a traditional practice that has been used for centuries. Companion planting can be used to improve soil health, attract beneficial insects, and repel pests. This publication discusses the scientific and traditional basis for companion planting associations including soil mixing, weed suppression, physical spatial resistance, and other mechanisms. It provides a companion planting chart for common herbs, vegetables, and flowers, as well as a listing of literature resources. An appendix provides information on the Three Sisters, a traditional Native American companion planting practice.

Botanical Self-Defense
 Plants have evolved various chemical and physical defenses to protect themselves from herbivores. Some plants have evolved chemical defenses that are toxic to herbivores. Other plants have evolved physical defenses such as thorns, spines, and trichomes. This publication discusses the scientific and traditional basis for botanical pesticides including soil mixing, weed suppression, physical spatial resistance, and other mechanisms. It provides a botanical pesticide chart for common herbs, vegetables, and flowers, as well as a listing of literature resources. An appendix provides information on the Three Sisters, a traditional Native American companion planting practice.

Spider Mites
 Spider mites are a common pest of many plants. They are tiny, eight-legged arachnids that feed on plant tissue. Spider mites can cause significant damage to plants, especially in hot, dry conditions. This publication discusses the scientific and traditional basis for spider mite control, including companion planting and botanical pesticides. It provides a list of plants that are resistant to spider mites and a list of botanical pesticides that are effective against spider mites.

Companion Planting
 Companion planting is a practice of growing different plants together to benefit them or to repel pests. It is a traditional practice that has been used for centuries. Companion planting can be used to improve soil health, attract beneficial insects, and repel pests. This publication discusses the scientific and traditional basis for companion planting associations including soil mixing, weed suppression, physical spatial resistance, and other mechanisms. It provides a companion planting chart for common herbs, vegetables, and flowers, as well as a listing of literature resources. An appendix provides information on the Three Sisters, a traditional Native American companion planting practice.

Botanical Pesticides
 Botanical pesticides are pesticides derived from natural sources. They are often used as alternatives to synthetic pesticides. This publication discusses the scientific and traditional basis for botanical pesticides including soil mixing, weed suppression, physical spatial resistance, and other mechanisms. It provides a botanical pesticide chart for common herbs, vegetables, and flowers, as well as a listing of literature resources. An appendix provides information on the Three Sisters, a traditional Native American companion planting practice.

Spider Mites
 Spider mites are a common pest of many plants. They are tiny, eight-legged arachnids that feed on plant tissue. Spider mites can cause significant damage to plants, especially in hot, dry conditions. This publication discusses the scientific and traditional basis for spider mite control, including companion planting and botanical pesticides. It provides a list of plants that are resistant to spider mites and a list of botanical pesticides that are effective against spider mites.

www.attra.ncat.org Page 1

ATTRA has a database of low-risk pesticides, including some botanicals, at <https://attra.ncat.org/attra-pub/biorationals>. For more information, consult the ATTRA publication Companion Planting & Botanical Pesticides: Concepts & Resources at attra.ncat.org/attra-pub/summaries/summary.php?pub=72.

This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2013-51106-20970.

Spider Mites—Botanical Control Formulations

By Justin Duncan, NCAT Agriculture Specialist • Published August 2016 ©NCAT IP522 • Slot 548



Produced by the National Center for Appropriate Technology
www.ncat.org • 1-800-275-6228 (1-800-ASK-NCAT)
 (Parent organization to the ATTRA Project, www.attra.ncat.org)